



Functional Unit

Scenario name:	PSB
Product:	Marley Ballpen
Primary Material:	
Scope:	Cradle - Gate

Total Results

 GHG
(kg Co2-eq)

0.1005

 Water Use
(m3 world eq. deprived)

0.0420

 Land Use
(points)

0.2596

Main Environmental Indicators

 GHG
Kg Co2-eq

The increase in the average global temperatures for 100 years as result of greenhouse gas (GHG) emissions. The greatest contributor is generally the combustion of fossil fuels such as coal, oil, and natural gas. Divided into 3 subcategories based on the emission source: (1) fossil resources, (2) bio-based resources, and (3) land use change.

Results:	0.1005	Kg Co2-eq
Benchmark:	0.0885	Kg Co2-eq
Savings %:	-13.5319	
Avoided:	-0.0120	Kg Co2-eq

 Water Use
m3 world eq. deprived

The withdrawal of water from lakes, rivers or groundwater can contribute to the 'depletion' of available water. The impact category considers the availability or scarcity of water in the regions where the activity takes place, if this information is known.

Results:	0.0420	m ³ world eq. deprived
Benchmark:	0.0407	m ³ world eq. deprived
Savings %:	-3.1937	
Avoided:	-0.0013	m ³ world eq. deprived

 Land Use
Points

Use and transformation of land for agriculture, roads, housing, mining or other purposes. The impacts can vary and include loss of species, of the organic matter content of soil, or loss of the soil itself (erosion). This is a composite indicator measuring impacts on four soil properties (biotic production, erosion resistance, groundwater regeneration and mechanical filtration).

Results:	0.2596	Points
Benchmark:	0.2258	Points
Savings %:	-14.9758	
Avoided:	-0.0338	Points