



Functional Unit

Scenario name:	UKX
Product:	Odyssey Pack Recycled Cotton Canvas Tote Bag- 17 L
Primary Material:	
Scope:	Cradle - Gate

Total Results

GHG (kg Co2-eq)	Water Use (m3 world eq. deprived)	Land Use (points)
10.0618	2.9011	30.6156

Main Environmental Indicators



The increase in the average global temperatures for 100 years as result of greenhouse gas (GHG) emissions. The greatest contributor is generally the combustion of fossil fuels such as coal, oil, and natural gas. Divided into 3 subcategories based on the emission source: (1) fossil resources, (2) bio-based resources, and (3) land use change.

Results:	10.0618	Kg Co2-eq
Benchmark:	10.3319	Kg Co2-eq
Savings %:	2.6135	
Avoided:	0.2701	Kg Co2-eq



The withdrawal of water from lakes, rivers or groundwater can contribute to the 'depletion' of available water. The impact category considers the availability or scarcity of water in the regions where the activity takes place, if this information is known.

Results:	2.9011	m ³ world eq. deprived
Benchmark:	10.3033	m ³ world eq. deprived
Savings %:	71.8430	
Avoided:	7.4022	m ³ world eq. deprived



Use and transformation of land for agriculture, roads, housing, mining or other purposes. The impacts can vary and include loss of species, of the organic matter content of soil, or loss of the soil itself (erosion). This is a composite indicator measuring impacts on four soil properties (biotic production, erosion resistance, groundwater regeneration and mechanical filtration).

Results:	30.6156	Points
Benchmark:	75.2847	Points
Savings %:	59.3335	
Avoided:	44.6691	Points