




Functional Unit

Scenario name:	ULA
Product:	Taiga rPET Nylon Zippered Waist Bag - 1.2 L
Primary Material:	
Scope:	Cradle - Gate

Total Results

 GHG (kg Co2-eq)	 Water Use (m3 world eq. deprived)	 Land Use (points)
4.0334	0.8332	9.2692

Main Environmental Indicators



The increase in the average global temperatures for 100 years as result of greenhouse gas (GHG) emissions. The greatest contributor is generally the combustion of fossil fuels such as coal, oil, and natural gas. Divided into 3 subcategories based on the emission source: (1) fossil resources, (2) bio-based resources, and (3) land use change.

Results:	4.0334	Kg Co2-eq
Benchmark:	4.1128	Kg Co2-eq
Savings %:	1.9744	
Avoided:	0.0794	Kg Co2-eq



The withdrawal of water from lakes, rivers or groundwater can contribute to the 'depletion' of available water. The impact category considers the availability or scarcity of water in the regions where the activity takes place, if this information is known.

Results:	0.8332	m ³ world eq. deprived
Benchmark:	0.8660	m ³ world eq. deprived
Savings %:	3.8594	
Avoided:	0.0328	m ³ world eq. deprived



Use and transformation of land for agriculture, roads, housing, mining or other purposes. The impacts can vary and include loss of species, of the organic matter content of soil, or loss of the soil itself (erosion). This is a composite indicator measuring impacts on four soil properties (biotic production, erosion resistance, groundwater regeneration and mechanical filtration).

Results:	9.2692	Points
Benchmark:	9.0238	Points
Savings %:	-2.7196	
Avoided:	-0.2454	Points